

## Surgical Pre-Operative Medications

**\*\*Please *start day before* Surgery\*\***

- Medrol Dose Pack / or Methyl Trednisolone

A medication to help suppress bruising and swelling. Take as directed until finished (6 days). \* Directions located under each tab in packet\*.

**\*\*Please *start day of* surgery\*\***

- Amoxicillin 500 mg. / Other Antibiotic: \_\_\_\_\_

Take one (1) tablet 4 hours before surgery

Take one (1) tablet at time of surgery

Following your surgery, take one (1) tablet every 6 hours (in your waking hours) until finished (10 days)

\*\*If you are a Pre-medication Patient, you will follow your pre-medication instructions.

- Motrin 800mg. / or Ibuprofen 800mg.

Take one (1) tablet 4 hours before surgery

Take one (1) tablet at time of surgery

Following surgery take one (1) tablet every 4-6 hours as needed for pain. Stay on this medication for 2-3 days to keep inflammation down.

NORCO will also be prescribed for pain is more intense. Refer to homecare instructions that will be given on day of surgery.

Other Pain Medication: \_\_\_\_\_

**\*24 hours after surgery\***

Peridex Antibacterial Rinse / Perio Guard or Chlorhexidine

Use as directed by Doctor or Assistant. (Refer to homecare instructions).

**\*\*YOU MUST HAVE AN ADULT ACCOMPANY YOU TO AND FROM SURGERY WHEN UNDER SEDATION\*\***

### **Sedation Instructions for the day of Surgery**

Halcion.25mg / or Triazolam

Take one (1) tablet 1 hour before surgery

Take one (1) tablet at time of surgery ***if needed***

**OR**

Valium 10mg. / or Diazepan

Take one (1) tablet 1 hour before surgery

Take one (1) tablet at time of surgery ***if needed***

**You may eat a light meal before your surgery.**

**Please fill out this form for the surgical staff the day of your surgery.**

Be sure to start Medrol pack the day BEFORE surgery, and continue following instructions on packet.

**Day Of Surgery:**

Your surgery is scheduled at \_\_\_\_\_

**4 Hours Before Surgery** (\_\_\_\_\_:\_\_\_\_\_) )

Antibiotic(\_\_\_\_\_) Taken: Y / N

Motrin (Ibuprophen) Taken: Y / N

**1 Hour Before Surgery** (\_\_\_\_\_:\_\_\_\_\_) )

Halcion or Valium Taken: Y / N

**At Time Of Surgery** –Surgical Staff to administer this dose of medication.

Time given to patient (\_\_\_\_\_:\_\_\_\_\_) )

Antibiotic(\_\_\_\_\_) Taken: Y / N

Motrin (Ibuprophen) Taken: Y / N

IF NEEDED:

Halcion or Valium Taken: Y / N

**For further medication directions, please refer to homecare instructions.**